



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$\lambda\lambda 463$ and 467 are possibly the titanium flutings at $\lambda\lambda 4626$ and 4668 . The latter appears as a bright line on one of my plates.

The plate obtained on the nights of July 11th and 12th of this year, exposure 10.2 hours, gives from the absorption lines a radial velocity of -30 km. The displacement of some of the bright lines is shown in the following table:

λ (measured)	Identification	Displacement	Weight
6565.2	H α	+2.4 A	½
4800.0	H β	-1.3 "	2
4686.0	4685.8 (nebular)	+0.2 "	1
4660.6	4658.2 (nebular)	+2.4 "	1
4365.1	4363.2 (nebular)	+1.9 "	1
4348.9	4347.0 (P Cygni?)	+1.9 "	½

Lick Observatory, September, 1921.

KNUT LUNDMARK.

NOTE ON THE SPECTRUM OF COMPONENT C IN THE TRIPLE SYSTEM
 σ^2 ERIDANI

From a plate taken 1921 October 7 with the 36-inch refractor, light 1-prism spectrograph, and 6-inch camera, the spectrum of the fainter component (C) of the double companion BC to σ^2 Eridani (=Burnham's G. C. No. 2109: BC: 1921.69, p. a. = 18° , dist. = $3\frac{1}{2}''$, B = $9^m.1$, C = $10^m.8$, visual) may be provisionally classified as Md; H β , at least, is bright. The negative is weak, having been obtained thru clouds, but a stronger effective exposure will be secured immediately. On the assumption that the parallax of the triple star is $o''.196$, the absolute magnitude of C is +12.3. According to Dr. Adams¹, Ao is the spectral class of component B.

FREDERICK C. LEONARD.

Lick Observatory,
Mount Hamilton, Cal., 1921 Oct. 8.

RETURN OF ASTRONOMICAL INSTRUMENTS TO CHINA.

A letter just received from Professor Chang, Acting Director of the Central Observatory, Peking, China, brings the interesting news that the five bronze astronomical instruments purloined from the Peking Observatory at the time of the Boxer Rebellion by the officers of the German punitive forces in China, shipped to Germany and mounted on the lawn of the Kaiser's palace at Potsdam

¹Publ. A. S. P. 26 (1914), 198.